

You have a

Soft tissue foot injury

This is a sprain to your foot.

Healing:

It normally takes 6 weeks for this to heal.

Smoking will slow down your healing. We would advise that you stop smoking while your fracture heals. Talk to your GP or go to www.smokefree.nhs.uk for more information.

Pain and swelling:

You may have foot pain and swelling for 3-6 months after your injury. Swelling is often worse at the end of the day.

Taking pain medication, elevating your foot and using ice or cold packs will help. More information is on the next page.



Walking and your boot:

The boot protects your foot and will make you more comfortable. Wear the boot when you are standing and walking. You can take it off at night and at rest. You need to wear the boot for 2 weeks after your injury. Please inform us if you are diabetic; you may require a specialist boot.

You are allowed to put weight through your foot. You may find it easier to use crutches in the early stages.



Exercises:

It is important to start exercises as soon as possible. Instructions are on the next page.

Follow up:

A follow up appointment is not normally needed for this injury. If you still have significant pain and swelling after 3 months, then please contact the Virtual Fracture Clinic team.

Any questions:

If you are concerned about your symptoms, are unable to follow this rehabilitation plan or have pain other than at the site of your injury please contact the Virtual Fracture Clinic team.

Remember to wear your boot whenever standing and walking. You can remove the boot when resting, at night or to wash and dress. Wear a long sock in your boot.

Using a cold pack will help with your pain and swelling. You can use an ice pack or bag of frozen peas wrapped in a damp towel. Put this on your foot for up to 15 minutes every few hours. Make sure the ice is not in direct contact with your skin.

Try to rest your foot, especially in the first 24-72 hours. Raise your foot on a stool or cushions so that it is above the level of your hip. This will help to reduce your swelling.


Exercises:

Early movement of the ankle and foot is important to promote circulation and reduce the risk of developing a Deep Vein Thrombosis (blood clot).

Do these exercises 3-4 times a day. Start straight away, you do not need to push into pain.



1. Point your foot up and down. Repeat this 10 times.



2. With your heels together, move your toes apart to turn the foot outwards. Repeat this 10 times. Do this movement gently within comfort.



3. Make gentle circles with your foot in one direction and then the other direction. Repeat this 10 times.

It is now 2 weeks since your injury and you can start to wean out of your boot. Stop using it around your own home first and continue wearing it for longer distances. You should aim to be fully weaned out of your boot by 6 weeks after injury at the latest.

It is normal to still have mild discomfort and swelling. This may continue for 3-6 months.

Activity and Exercise

Keep doing your exercises until you have full movement of your foot.

Gradually increase your level of activity. You should avoid impact activity for 3 months. This includes running, jumping and dancing.

I am struggling with my boot. What do I do?

The boot has a thicker sole; this can make you feel uneven. We would recommend choosing a supportive shoe or trainer with a firm sole for your uninjured foot. If you are struggling to wear out of your boot please contact us.

If you need more advice contact the Virtual Fracture Clinic.

I am diabetic, does this change things?

If you are diabetic please contact us to discuss your boot. This is particularly important if you have problems with your skin. We may provide you with a specialist diabetic boot.

When can I start driving?

You can return to driving when:

- You are no longer using your boot,
- You can walk comfortably
- You can perform an emergency stop pain free.

Always test your ability to drive in a safe environment first.

How can I get a certificate for work?

You can get a fitness for work statement from your GP.

What do I do with my boot and crutches when I no longer need them?

We are not able to use boots again. These should not be returned to the hospital.

Crutches can be returned to the Fracture Clinic or A&E.

How do I contact the Virtual Fracture Clinic?

Call 01273 696955 extension 63428.

Email uhsussex.fracturecare@nhs.net